

# All-Trans Retinoic Acid Inhibits Fluctuations in Intracellular $\text{Ca}^{2+}$ Resulting from Changes in Extracellular $\text{Ca}^{2+}$

James Varani, Bill Burmeister,  
Patricia Perone, Michael Bleavins,  
and Kent J. Johnson

*From the Department of Pathology, The University of Michigan, and the Department of Pathology and Experimental Toxicology, Parke-Davis Pharmaceutical Research, Division of the Warner-Lambert Co., Ann Arbor, Michigan*

**Previous studies have shown that all-trans retinoic acid (RA) preserves fibroblast viability and stimulates their proliferation, in part, by reducing the extracellular  $\text{Ca}^{2+}$  requirement (Am J Pathol 1990, 130:1275). Based on this observation, we have in the present study examined the effects of RA on  $\text{Ca}^{2+}$  mobilization in human dermal fibroblasts. For these studies we used the  $\text{Ca}^{2+}$ -binding dyes, Fluo-3 and Indo-1. Using fluorescence of Fluo-3-loaded cells or Indo-1-loaded cells as indicators of intracellular free  $\text{Ca}^{2+}$ , we observed that treatment of the cells with RA did not, by itself, alter the concentration of intracellular  $\text{Ca}^{2+}$ . Nor did it interfere with the rapid, transient rise in intracellular  $\text{Ca}^{2+}$  induced by treatment with ionomycin. However, treatment of the cells with RA prevented re-equilibration of intracellular  $\text{Ca}^{2+}$  when the cells were initially equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) culture medium and then switched to high  $\text{Ca}^{2+}$  (1.4 mmol/L) medium or when cells were first equilibrated in high  $\text{Ca}^{2+}$  medium and then switched to low  $\text{Ca}^{2+}$  medium. This effect of RA could be seen within seconds after treatment and the effect was observed 1 day after treatment (longest time point examined). The effect was concentration dependent and concentrations of RA that modulated  $\text{Ca}^{2+}$  re-equilibration (0.3 to 3.0  $\mu\text{mol/L}$ ) were the same as those that have previously been shown to promote fibroblast survival and growth. A biologically inactive retinoid did not have this effect. Specificity of the response was**

**suggested by the finding that concentrations of RA that modulated  $\text{Ca}^{2+}$  movement had no effect on  $\text{Ba}^{2+}$  transport. These data suggest that RA prevents re-equilibration of intracellular  $\text{Ca}^{2+}$  in human dermal fibroblasts by interfering with  $\text{Ca}^{2+}$  movement across the plasma membrane. (Am J Pathol 1995, 147:718–727)**

Topical application of all-trans retinoic acid (RA) improves the appearance of sun-damaged skin,<sup>1,2</sup> and recent studies suggest that RA may have a beneficial effect in natural aging as well.<sup>3</sup> How retinoids counteract the effects of aging in skin is not fully understood, but changes in both the epidermis and dermis occur. The epidermis of retinoid-treated skin is thickened, which is caused, in part, by increased keratinocyte proliferation and increased deposition of water-retaining glycosaminoglycans. In the dermis, one sees evidence of fibroblast activation and new extracellular matrix synthesis. Recent studies from our laboratory with an organ culture model suggest that retinoid activation of the dermis is critical, not only for preservation of the dermis itself but also for keratinocyte proliferation and epidermal thickening.<sup>4–7</sup>

The molecular and biochemical mechanisms underlying the therapeutic effects of retinoids have not been delineated. Past studies have shown that retinoids activate transcription of collagen I genes to enhance collagen synthesis.<sup>8,9</sup> Additionally, retinoids can decrease production of collagen-degrading enzymes.<sup>10</sup> Together, these specific effects on collagen metabolism could account for the deposition of new collagen in the zone immediately below the dermo-epidermal basement membrane in retinoid-treated skin.<sup>11</sup> Studies in

---

Supported in part by grant CA60958 from the US Public Health Service.

Accepted for publication May 15, 1995.

Address reprint requests to Dr. James Varani, Department of Pathology, The University of Michigan Medical School, 1301 Catherine Rd., Box 0602, Ann Arbor, MI 48109.

rodent models of photodamage support the concept that deposition of new collagen directly underlies clinical improvement.<sup>12-14</sup> The question is whether these changes in collagen metabolism reflect retinoid effects on specific genes or whether the changes reflect a more generalized enhancement of fibroblast function. Our own studies suggest that the retinoid influence on fibroblast function is a generalized effect and tied in some manner to modulation of  $\text{Ca}^{2+}$ -regulated processes. This is based on the finding that human dermal fibroblasts normally require an extracellular  $\text{Ca}^{2+}$  concentration of approximately 1 mmol/L or higher for survival and growth<sup>15-20</sup> and that, whereas a variety of growth factors including epidermal growth factor and insulin-like growth factor-1 cannot replace this  $\text{Ca}^{2+}$  requirement, the addition of RA allows for survival and growth at extracellular  $\text{Ca}^{2+}$  concentrations as low as 0.15 mmol/L.<sup>4,21</sup> How RA functions to maintain fibroblast survival and growth at a suboptimal level of extracellular  $\text{Ca}^{2+}$  is not understood. To begin addressing this question, we have, in the present study, examined the effects of RA on intracellular  $\text{Ca}^{2+}$  levels in human dermal fibroblasts. Automated laser fluorescence microscopy was used in conjunction with the fluorescent  $\text{Ca}^{2+}$ -binding dyes Fluo-3 and Indo-1 for these studies, allowing measurements to be made on individual cells while maintaining the cells in monolayer culture.

## Materials and Methods

### Culture Media and Reagents

Keratinocyte growth medium (KGM) was used as culture medium for these experiments. KGM is a serum-free modification of MCDB-153 medium and is supplemented with a number of growth factors including epidermal growth factor, insulin, hydrocortisone, and pituitary extract (Clonetics, San Diego, CA). The  $\text{Ca}^{2+}$  concentration of KGM is 0.15 mmol/L. Dulbecco's modified minimal essential medium of Eagle supplemented with Earle's salts, nonessential amino acids, and 10% fetal bovine serum (DMEM-FBS) was also used. The medium and additives were obtained from GIBCO (Grand Island, NY) and the fetal bovine serum was obtained from Hyclone Laboratories (Logan, UT).

RA was obtained from the R. W. Johnson Pharmaceutical Research Institute (Raritan, NJ) and prepared as a 20-mg/ml solution in dimethyl sulfoxide (DMSO). The stock solution was stored at  $-20^{\circ}\text{C}$  in the dark until use. At the time of use, it was diluted directly in KGM and added to the cultures. In cultures treated with 3  $\mu\text{mol/L}$  RA (highest concentra-


tion used), the final concentration of DMSO in the culture medium was 0.005%. This amount of DMSO had no detectable effect by itself. In certain experiments, the biologically inactive retinoid, M-[(E)-2-(5, 6, 7, 8-tetrahydro-5, 5, 8, 8-tetramethyl-2-naphthalenyl-1-propenyl)] benzoic acid (mc-TTNPB), was used in place of RA. In other experiments, 1,25-dihydroxyvitamin  $\text{D}_3$  (vitamin D) was used. Both were obtained from Hoffmann-La Roche (Nutley, NJ). Both mc-TTNPB and vitamin D were prepared in DMSO and handled as RA. Neither of these agents supports fibroblast survival and growth in low  $\text{Ca}^{2+}$  culture medium.<sup>4,22</sup>

### Cells

Fibroblasts were obtained from neonatal foreskin and cultured in DMEM-FBS as previously described.<sup>23</sup> Keratinocytes were obtained from neonatal foreskin by the method of Liu and Karasek<sup>24</sup> and grown in KGM. Both fibroblasts and keratinocytes were grown at  $37^{\circ}\text{C}$  in an atmosphere of 5%  $\text{CO}_2$  and subcultured as required. Cells were routinely used at passages 2 to 5. For use in  $\text{Ca}^{2+}$  measurement studies, the cells were plated at  $4 \times 10^4$  cells per well in a 24-well culture dish. KGM was used as plating medium for keratinocytes and DMEM-FBS was used as plating medium for fibroblasts. After allowing the cells to attach and spread, cells were washed twice in KGM (containing either the normal extracellular  $\text{Ca}^{2+}$  concentration of 0.15 mmol/L or supplemented with extracellular  $\text{CaCl}_2$  to bring the final concentration to 1.4 mmol/L  $\text{Ca}^{2+}$ ). At this point, the cells were ready for use.

### $\text{Ca}^{2+}$ Measurement Assays

For fluorescent measurements, cells were first incubated with the acetoxymethyl ester form of Fluo-3 (5  $\mu\text{mol/L}$ ) by incubation for 30 minutes at room temperature in KGM (0.15 or 1.4 mmol/L  $\text{Ca}^{2+}$ ). After 30 minutes, the cells were washed twice in KGM and fresh KGM containing either 0.15 mmol/L or 1.4 mmol/L  $\text{Ca}^{2+}$  was added. The 24-well culture dish was then mounted on the stage of an ACAS-570 interactive confocal laser cytometer (Meridian Instruments, Okemos, MI). The labeled cells were then excited with 488-nm light from an argon laser, and the emission greater than 515 nm was detected with a photomultiplier tube and digitized. Once control fluorescence values had stabilized, the desired reagents were added directly to the wells. Fluorescence values were then monitored for the remainder of the experiment. Ionomycin (0.5  $\mu\text{mol/L}$ ) was



**Figure 1.** Fluo-3 fluorescence emission profile of human dermal fibroblasts. **A:** Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM.  $\text{CaCl}_2$  was added at the point indicated to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. **B:** Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  RA. **C:** Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  RA and  $\text{CaCl}_2$  to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. Each panel represents cells from a single well in one experiment. Quadruplicate wells were routinely assayed for each condition in each experiment. The experiment was repeated with fibroblasts isolated from six different foreskins with virtually identical results.

added at the conclusion of each experiment. This agent is a  $\text{Ca}^{2+}$  ionophore. It allows  $\text{Ca}^{2+}$  influx into the cells and is used to generate the maximal fluorescent signal. (Methodological details can be found in Meridian application note E-2.)


In additional studies, cells in eight-chamber glass slides were loaded with the acetoxymethyl ester form of the  $\text{Ca}^{2+}$ -sensitive dye Indo-1 (Indo-1-AM). This was done by incubating the cells for 1 hour at 37°C with 1  $\mu\text{mol/L}$  Indo-1-AM in KGM (0.15 mmol/L  $\text{Ca}^{2+}$ ). After the 1-hour incubation, the cells were washed two times in KGM and fresh KGM containing 0.15 mmol/L  $\text{Ca}^{2+}$  was added. The slides were then mounted on the stage of the ACAS-570 interactive confocal laser cytometer. The labeled cells were excited with 355-nm light and two emission peaks at 404 nm and 485 nm were detected with a photomultiplier tube and digitized. (The 404-nm peak increases in conjunction with increasing  $\text{Ca}^{2+}$  concentration whereas the 485-nm peak decreases with increasing  $\text{Ca}^{2+}$  concentration.) Once control fluorescence values had stabilized, the desired reagents were added directly to the wells. Fluorescence values were then monitored for the remainder of the experiment. Ionomycin (0.5  $\mu\text{mol/L}$ ) and EGTA (10 mmol/L) were added in sequence at the conclusion of each experiment. These two agents were used to generate maximal and minimal fluorescent signals. Intracellular  $\text{Ca}^{2+}$  levels were then calculated by using the relationship  $\text{Ca}^{2+} = K_d S ([R - R_{\min}]/[R_{\max} - R])$  as described by Grynkiewicz et al.<sup>25</sup> The published value for the effective dissociation constant ( $K_d$ ) of the dye (ie, 250 nmol/L)<sup>25</sup> was used in the calculation.  $R$  is the ratio of the fluorescence emission at 404 nm and 485 nm and  $R_{\min}$  and  $R_{\max}$  are the ratios in the presence of EGTA and ionomycin. In addition to calculating intracellular  $\text{Ca}^{2+}$  values, a  $\text{Ca}^{2+}$  standard curve was generated and the intracellular  $\text{Ca}^{2+}$  values determined by direct comparison with the standard curve. (Methodological details can be found in Meridian Instruments application note entitled Intracellular Calcium Quantitation Using the ACAS 570.)

## Results

### Intracellular Fluorescence Levels in Fibroblasts

#### Effects of RA on Cells Equilibrated in Low $\text{Ca}^{2+}$ (0.15 mmol/L) Culture Medium and Shifted to High $\text{Ca}^{2+}$ (1.4 mmol/L) Medium

In the first series of experiments, Fluo-3-treated fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and analyzed for fluorescence emission. When a stable emission pattern was obtained,  $\text{CaCl}_2$  was introduced into the well to bring the extracellular  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. This resulted in a



**Figure 2.** Indo-1 fluorescence emission profile of human dermal fibroblasts. Indo-1-labeled fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM.  $\text{CaCl}_2$  was added at the point indicated to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. Ionomycin and EGTA were added at the points indicated. Dye-labeled cells were excited with 355-nm light and fluorescence emissions detected at 404 and 485 nm. Intracellular  $\text{Ca}^{2+}$  values were obtained by calculation using the previously published formula.<sup>25</sup> Inset: A calcium standard curve was generated by adding various amounts of  $\text{Ca}^{2+}$  to the assay buffer. The  $\text{Ca}^{2+}$ -containing buffers were incubated with the free acid of Indo-1. The buffers were excited with 355-nm light and fluorescence emissions detected at 404 and 485 nm. Intracellular  $\text{Ca}^{2+}$  values were obtained by direct comparison with the standard curve.


rapid increase in intracellular fluorescence, observed in virtually every cell. The increase in fluorescence was routinely in the range of 2.5- to 3-fold and the maximal change was seen within 1 to 2 minutes. Thereafter, fluorescence emission values decreased to an intermediate level and remained relatively constant. Figure 1A shows results from a typical experiment in which fluorescence emission of individual cells is plotted as a function of time. A disadvantage of using Fluo-3 to determine intracellular  $\text{Ca}^{2+}$  concentrations is that it produces a single-wavelength emission, and because of this, it is difficult to obtain absolute  $\text{Ca}^{2+}$  concentrations. Therefore, parallel experiments were conducted in which Indo-1 was used in place of Fluo-3 as the  $\text{Ca}^{2+}$ -binding dye. As in the Fluo-3 experiments, cells were equilibrated with KGM containing 0.15 mmol/L  $\text{Ca}^{2+}$  and then shifted to 1.4 mmol/L  $\text{Ca}^{2+}$ . A change in intracellular fluorescence was observed upon shifting (Figure 2) and from the fluorescence changes (ie, at 404 nm and 485 nm), the intracellular  $\text{Ca}^{2+}$  concentrations were calculated. In low  $\text{Ca}^{2+}$ -containing medium, the intracellular  $\text{Ca}^{2+}$  concentration was calculated to be 65 nM and this increased to 175 nM under conditions of high-extracellular  $\text{Ca}^{2+}$ .

Figure 1B shows the effects of RA on intracellular  $\text{Ca}^{2+}$  concentration. For this experiment, Fluo-3-containing cells were equilibrated with low  $\text{Ca}^{2+}$  KGM and then treated with 3  $\mu\text{mol/L}$  RA. This treatment resulted in no increase in the fluorescence emission pattern. If anything, there was a slight reduction in emission, suggestive of decreased intracellular  $\text{Ca}^{2+}$ . As part of the same experiment, dye-loaded fibroblasts were treated with 3  $\mu\text{mol/L}$  RA in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and immediately

shifted to 1.4 mmol/L  $\text{Ca}^{2+}$ . In contrast to what was seen in control cells, fluorescence emission levels did not rise in cells that were treated with RA along with the extracellular  $\text{Ca}^{2+}$  (Figure 1C). The data shown in Figure 1C are based on readings from individual cells in a single well. In each experiment, quadruplicate wells were used and no significant differences among the different wells were observed. The experiment was repeated with fibroblasts isolated from six different foreskins and virtually identical results were obtained with all six. Thus it appears that treatment of fibroblasts with RA not only does not result in an increase in intracellular  $\text{Ca}^{2+}$  concentration but actually retards the increase seen after addition of  $\text{Ca}^{2+}$  to the extracellular environment. In an additional experiment, Indo-1-loaded cells equilibrated in 0.15 mmol/L  $\text{Ca}^{2+}$  were exposed to 3  $\mu\text{mol/L}$  RA and 1.4 mmol/L  $\text{Ca}^{2+}$  concomitantly and fluorescence emission patterns were monitored. As expected, the presence of RA completely inhibited the increase in 404-nm fluorescence and the decrease in 485-nm fluorescence (not shown).

Although RA inhibited the rise in intracellular fluorescence resulting from the increase in extracellular  $\text{Ca}^{2+}$ , the rapid, transient rise in intracellular fluorescence that occurred after addition of 0.5  $\mu\text{mol/L}$  ionomycin was not inhibited by this treatment (Figure 1, B and C). The rapid, transient rise in intracellular  $\text{Ca}^{2+}$  concentration seen after ionomycin treatment is caused primarily by a release of  $\text{Ca}^{2+}$  from intracellular stores (endoplasmic reticulum and, perhaps, mitochondria<sup>26-28</sup>).

Figure 3 presents two-dimensional fluorescence emission images of Fluo-3-labeled cells (equilibrated in 0.15 mmol/L  $\text{Ca}^{2+}$ ) before and after the addition of




**Figure 3.** Two-dimensional fluorescence emission images of Fluo-3-labeled human dermal fibroblasts before and after the addition of 1.4 mmol/L  $\text{Ca}^{2+}$  (final concentration) alone or in conjunction with 3  $\mu\text{mol/L}$  RA. **A:** Control cells before addition of  $\text{Ca}^{2+}$ . **B:** Control cells 40 seconds after addition of  $\text{Ca}^{2+}$ . **C:** Control cells 6 minutes after addition of  $\text{Ca}^{2+}$ . **D:** Cells before addition of RA and  $\text{Ca}^{2+}$ . **E:** Cells 40 seconds after addition of RA and  $\text{Ca}^{2+}$ . **F:** Cells 6 minutes after addition of RA and  $\text{Ca}^{2+}$  and 40 seconds after addition of 0.5  $\mu\text{mol/L}$  ionomycin.

$\text{Ca}^{2+}$  to bring the extracellular concentration to 1.4 mmol/L  $\text{Ca}^{2+}$ . Control cells equilibrated in 0.15 mmol/L extracellular  $\text{Ca}^{2+}$  are shown in Figure 3, A–C. There is a rapid increase in fluorescence upon addition of  $\text{Ca}^{2+}$  (Figure 3B) with a subsequent decrease to an intermediate level (Figure 3C). In cells treated with 3  $\mu\text{mol/L}$  RA, there is no change in intracellular fluorescence upon addition of  $\text{Ca}^{2+}$  (Figure 3, D and E) although cells treated with RA show a dramatic increase in intracellular fluorescence upon treatment with ionomycin (Figure 3F).


In additional experiments, varying concentrations of RA (3.0, 1.5, 0.75, 0.3, 0.15, and 0.075  $\mu\text{mol/L}$ ) were used. Concentrations as low as 0.3  $\mu\text{mol/L}$  were sufficient to inhibit by greater than 90% the increase in fluorescence emission elicited by addition of 1.4 mmol/L  $\text{Ca}^{2+}$  to the culture medium. In some experiments, a final RA concentration of 0.15  $\mu\text{mol/L}$  was also inhibitory, although 0.075  $\mu\text{mol/L}$  was not. The dose-response data are of interest because the same concentrations of RA that inhibit  $\text{Ca}^{2+}$ -induced fluorescence changes have previously been shown to preserve fibroblast viability and induce growth under low  $\text{Ca}^{2+}$  conditions.<sup>21,22</sup> Time-response studies demonstrated that inhibition of the increased intracellular fluorescence in response to addition of  $\text{Ca}^{2+}$  was still observable 1 day after RA treatment (longest time point examined) (Figure 4). Finally, experiments were conducted in which the biologically inactive retinoid mc-TTNPB<sup>22</sup> was

added to Fluo-3-labeled fibroblasts (3  $\mu\text{mol/L}$ ) in place of RA.  $\text{Ca}^{2+}$  was immediately added to bring the final concentration to 1.4 mmol/L and the fluorescence emission pattern was monitored in the normal manner (Figure 5). Unlike RA, the biologically inactive retinoid did not inhibit the ensuing change in fluorescence emission pattern. Likewise, substitution of 1,25-dihydroxy vitamin D<sub>3</sub> (0.1–1  $\mu\text{mol/L}$ ) for RA did not inhibit the  $\text{Ca}^{2+}$ -induced increase in intracellular fluorescence (not shown).



**Figure 4.** Long-term effects of RA on Fluo-3 fluorescence in human dermal fibroblasts. Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated for one day with 3  $\mu\text{mol/L}$  RA.  $\text{CaCl}_2$  was then added at the point indicated to bring the  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. Results represents cells from a single well in one experiment. Quadruplicate wells were routinely assayed in each experiment. The experiment was repeated with fibroblasts isolated from three different foreskins with virtually identical results.





**Figure 5.** Effects of mc-TNPB on Fluo-3 fluorescence in human dermal fibroblasts. Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  mc-TNPB and  $\text{CaCl}_2$  to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. Results represent cells from a single well in one experiment. Quadruplicate wells were routinely assayed in each experiment. The experiment was repeated with fibroblasts isolated from three different fore-skins with virtually identical results.


#### Effects of RA on Cells Equilibrated in High $\text{Ca}^{2+}$ (1.4 mmol/L) Culture Medium and Shifted to Low $\text{Ca}^{2+}$ (0.15 mmol/L) Medium

Experiments were conducted in which Fluo-3-loaded fibroblasts were equilibrated in high  $\text{Ca}^{2+}$ -containing KGM and then downshifted to low  $\text{Ca}^{2+}$  medium. Downshift was accomplished either by rapidly removing the culture medium containing 1.4 mmol/L  $\text{Ca}^{2+}$  and replacing it with medium containing 0.15 mmol/L  $\text{Ca}^{2+}$  or by addition of 10 mmol/L EGTA to the high  $\text{Ca}^{2+}$ -containing medium. In either case, there was a rapid decrease in intracellular fluorescence. Figure 6A shows results of an experiment in which EGTA was added directly to the high  $\text{Ca}^{2+}$ -containing medium. A rapid decrease in intracellular fluorescence occurred. The effects of RA (3  $\mu\text{mol/L}$ ) on changes in intracellular fluorescence after EGTA addition are shown in Figure 6B. The decrease in fluorescence was inhibited in the RA-treated cells.

#### Effects of RA on Changes in Intracellular Fluorescence in Fibroblasts


##### Studies with Barium ( $\text{Ba}^{2+}$ )

$\text{Ba}^{2+}$  has been used in the past to study the process of divalent cation transport. Past studies by others have shown that when  $\text{Ba}^{2+}$  is added to culture medium, it enters cells through divalent cation channels but more slowly than  $\text{Ca}^{2+}$ . When  $\text{Ba}^{2+}$ -treated



**Figure 6.** Effects of RA on Fluo-3 fluorescence in human dermal fibroblasts. **A:** Fibroblasts were equilibrated in high  $\text{Ca}^{2+}$  (1.4 mmol/L) KGM. At the time indicated, 10 mmol/L EGTA (final concentration) was added. **B:** Fibroblasts were equilibrated in high  $\text{Ca}^{2+}$  (1.4 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  RA and 10 mmol/L EGTA. Each panel represents cells from a single well in one experiment. Quadruplicate wells were routinely assayed for each condition in each experiment. The experiment was repeated with fibroblasts isolated from four different fore-skins with virtually identical results.

cells are placed in a low divalent cation-containing buffer,  $\text{Ba}^{2+}$  is not released from the cells into the buffer.<sup>26-28</sup> This is thought to reflect a failure of  $\text{Ba}^{2+}$  to be removed through the same efflux processes that remove intracellular  $\text{Ca}^{2+}$ . In the present study, Fluo-3-loaded cells were equilibrated in low  $\text{Ca}^{2+}$  KGM and then treated with 1.4 mmol/L  $\text{Ba}^{2+}$ . Consistent with past findings by others,<sup>26-28</sup> intracellular fluorescence increased, but the rate of increase was slower than that seen after addition of the same amount of  $\text{Ca}^{2+}$  (Figure 7A; compare with Figure 1A). Interestingly, when cells equilibrated in low  $\text{Ca}^{2+}$ -containing medium were treated with RA along with  $\text{Ba}^{2+}$ , there was no inhibition of the increase in fluorescent signal (Figure 7B). In the next experi-



**Figure 7.** Effects of RA on Fluo-3 fluorescence in human dermal fibroblasts. **A:** Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM.  $\text{BaCl}_2$  was added at the point indicated to bring the final  $\text{Ba}^{2+}$  concentration to 1.4 mmol/L. **B:** Fibroblasts were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  RA and  $\text{BaCl}_2$  to bring the final  $\text{Ba}^{2+}$  concentration to 1.4 mmol/L. Each panel represents cells from a single well in one experiment. Quadruplicate wells were routinely assayed for each condition in each experiment. The experiment was repeated with fibroblasts isolated from two different foreskins with virtually identical results.

ment, Fluo-3-treated fibroblasts were incubated in KGM containing 0.15 mmol/L  $\text{Ca}^{2+}$  and 1.4 mmol/L  $\text{Ba}^{2+}$ . Subsequently, the culture medium was rapidly replaced with KGM containing the same amount of  $\text{Ca}^{2+}$  but without  $\text{Ba}^{2+}$ . With this manipulation, the fluorescence emission pattern remained unchanged (not shown). It also remained unchanged when RA was included in the  $\text{Ba}^{2+}$ -free culture medium (not shown). Taken together, these data suggest that RA does not influence  $\text{Ba}^{2+}$  movement into or out of dermal fibroblasts.

### Intracellular Fluorescence Levels in Keratinocytes


#### Effects of RA on Cells Equilibrated in Low $\text{Ca}^{2+}$ (0.15 mmol/L) Culture Medium and Shifted to High $\text{Ca}^{2+}$ (1.4 mmol/L) Medium

In a final set of experiments, neonatal foreskin keratinocytes were examined in place of fibroblasts. For these experiments, keratinocytes were labeled with Fluo-3 and equilibrated in KGM containing 0.15 mmol/L  $\text{Ca}^{2+}$ . Fluorescence emission patterns were monitored in the normal manner. One group of cells was then treated with 1.4 mmol/L  $\text{Ca}^{2+}$  (final concentration) and a parallel group was treated with 3  $\mu\text{mol/L}$  RA and 1.4 mmol/L  $\text{Ca}^{2+}$ . Fluorescence emission patterns were again assessed in the normal manner. In contrast to what was observed with fibroblasts, the addition of  $\text{Ca}^{2+}$  to the extracellular buffer

(either with or without RA) did not result in a fluorescence change over the period of observation (400 seconds; Figure 8). Although keratinocytes did not demonstrate a rapid response to changes in extracellular  $\text{Ca}^{2+}$  concentration, intracellular fluorescence increased in these cells after addition of ionomycin, just as it did in fibroblasts (Figure 8).

### Discussion

Studies in our laboratory are directed toward understanding how retinoids influence the behavior of human dermal fibroblasts. Our past studies have shown that RA preserves viability and stimulates growth of fibroblasts in both monolayer culture and organ culture by reducing their requirement for extracellular  $\text{Ca}^{2+}$ .<sup>4,21</sup> Growth factors such as epidermal growth factor and insulin-like growth factor-1 do not mimic RA in this regard,<sup>6</sup> although studies by Betsholtz and Westermarck<sup>29</sup> showed that platelet-derived growth factor induced 3T3 fibroblast proliferation, in part, by lowering the extracellular  $\text{Ca}^{2+}$  requirement. A possible explanation for these findings is that RA acts to stimulate an increase in the intracellular  $\text{Ca}^{2+}$  concentration and thereby mimic the effects of having an adequate amount of extracellular  $\text{Ca}^{2+}$ . A rapid transient rise in intracellular  $\text{Ca}^{2+}$  concentration is seen in a number of different cell types upon stimulation with a variety of agonists<sup>26–28,30–33</sup> and is considered to be a critical



**Figure 8.** Effects of RA on Fluo-3 fluorescence in human epidermal keratinocytes. **A:** Keratinocytes were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM.  $\text{CaCl}_2$  was added at the point indicated to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. **B:** Keratinocytes were equilibrated in low  $\text{Ca}^{2+}$  (0.15 mmol/L) KGM and treated at the point indicated with 3  $\mu\text{mol/L}$  RA and  $\text{CaCl}_2$  to bring the final  $\text{Ca}^{2+}$  concentration to 1.4 mmol/L. Each panel represents cells from a single well in one experiment. Quadruplicate wells were routinely assayed for each condition in each experiment. The experiment was repeated with keratinocytes isolated from three different foreskins with virtually identical results.

event in many stimulus-coupled responses.<sup>34</sup> The findings presented here suggest, however, that this is not the case with RA. Using intracellular fluorescence in fibroblasts loaded with the  $\text{Ca}^{2+}$ -binding dye Fluo-3 as an indicator of intracellular  $\text{Ca}^{2+}$  concentration, we found no evidence for an alteration in intracellular  $\text{Ca}^{2+}$  concentration in response to RA treatment alone. Thus, we reject one of the principal hypotheses on which this study was originally based.

Although RA treatment by itself did not increase intracellular free  $\text{Ca}^{2+}$  levels, RA modulated the cellular response to changes in extracellular  $\text{Ca}^{2+}$  concentration. This was shown when RA-treated cells and control cells were either equilibrated in low  $\text{Ca}^{2+}$  culture medium and then shifted to high  $\text{Ca}^{2+}$  medium or equilibrated in high  $\text{Ca}^{2+}$  medium and then downshifted to low  $\text{Ca}^{2+}$  medium. In both cases, the intracellular  $\text{Ca}^{2+}$  level (as indicated by changes in intracellular fluorescence) rapidly re-equilibrated in control cells, whereas re-equilibration was inhibited in the RA-treated cells. In contrast, RA did not interfere with the rapid, transient intracellular  $\text{Ca}^{2+}$  changes induced by ionomycin. Our suggestion based on these findings is that RA modulates cell membrane properties in some manner to influence unstimulated  $\text{Ca}^{2+}$  movement across the membrane but does not interfere with processes that release  $\text{Ca}^{2+}$  from stored pools within the cell.

How RA acts to modulate  $\text{Ca}^{2+}$  movement into and out of cells is not known at this time.  $\text{Ca}^{2+}$

uptake by cells occurs through a number of mechanisms. Both voltage-gated channels and receptor-gated channels have been reported in mesenchymal cells.<sup>35–37</sup> Although not well characterized in fibroblasts, these channels presumably do not function in the absence of electrical or ligand stimulation. As RA prevented the movement of  $\text{Ca}^{2+}$  across the membrane in the absence of stimulation, it is unlikely that it acts by specifically inhibiting either receptor-gated or voltage-gated processes. In addition to these channels, there are pathways that allow divalent cations entry into cells in the absence of stimulation. Some of these are specific for  $\text{Ca}^{2+}$  whereas others are permeable to a number of divalent cations including  $\text{Ba}^{2+}$ .<sup>26–28</sup> We suggest that it is one or more of these pathways that is influenced by RA. The fact that we did not see a similar retinoid modulation of intracellular fluorescence when  $\text{Ba}^{2+}$  was used in place of  $\text{Ca}^{2+}$  suggests some specificity for  $\text{Ca}^{2+}$ .

These observations offer insight into how RA may function at the cellular level. Recent studies have focused on the ability of retinoids to influence gene transcription<sup>38,39</sup> after binding to nuclear receptors.<sup>40</sup> The present data do not rule out a role for gene transcription in retinoid action. They suggest, however, that other mechanisms may also be important. The rapidity with which effects on  $\text{Ca}^{2+}$ -induced fluorescent changes were seen would rule out synthesis of new mRNA and protein in this process. Even though induction of gene transcription is



not likely to underlie the RA effect on  $\text{Ca}^{2+}$  movement, this retinoid effect may still result from binding to specific receptor moieties rather than from some global membrane effect. This is suggested by the findings with the synthetic retinoid, mc-TTNPB. mc-TTNPB has detergent-like properties similar to RA<sup>41</sup> and is similar to RA in inducing fibroblast lysis at high concentrations.<sup>42</sup> Yet this agent does not stimulate fibroblast survival or growth.<sup>22</sup> Likewise, it does not modulate  $\text{Ca}^{2+}$  movement across the cell membrane (this report). Furthermore, mc-TTNPB is unlike RA and other biologically active retinoids in that it does not bind retinoid receptors.<sup>43</sup>

Whether RA-induced inhibition of  $\text{Ca}^{2+}$  movement across the fibroblast membrane plays a role in the overall mechanism by which retinoids influence aging skin is yet to be determined. However, it may. The concentrations of RA that interfere with  $\text{Ca}^{2+}$  movement across the fibroblast membrane (0.3 to 3.0  $\mu\text{mol/L}$ ) are the same as concentrations that stimulate fibroblast proliferation under low  $\text{Ca}^{2+}$  conditions<sup>21,22</sup> and that preserve skin structure and function in organ culture.<sup>5-7</sup> Most importantly, the same concentrations of RA are found in the viable portion of the skin after topical application of therapeutic amounts of RA.<sup>44</sup>

## References

- Kligman AM, Grove GL, Hirose R, Leyden JJ: Topical tretinoin for photoaged skin. *J Am Acad Dermatol* 1986, 15:836-859
- Weiss JS, Ellis CN, Headington JT, Tincoff T, Hamilton TA, Voorhees JJ: Topical tretinoin improves photoaged skin: a double-blind, vehicle-controlled study. *JAMA* 1988, 259:527-532
- Kligman AM, Dogadkina D, Lavker RM: Effects of topical tretinoin on non-sun-exposed skin of the elderly. *J Am Acad Dermatol* 1993, 29:25-33
- Varani J, Perone P, Griffiths CEM, Inman DR, Fligiel SEG, Voorhees JJ: All-*trans* retinoic acid (RA) stimulates events in organ-cultured skin that underlie repair. *J Clin Invest* 1994, 94:1747-1756
- Varani J, Fligiel SEG, Schuger L, Perone P, Inman DR, Griffiths CEM, Voorhees JJ: Effects of all-*trans* retinoic acid, and  $\text{Ca}^{++}$  on human skin in organ culture. *Am J Pathol* 1993, 142:189-198
- Varani J, Fligiel SEG, Perone P, Inman DR, Voorhees JJ: Effects of sodium lauryl sulfate on human skin in organ culture: comparison with all-*trans* retinoic acid and epidermal growth factor. *Dermatology* 1993, 187:19-25
- Varani J, Larson BK, Perone R, Inman DR, Fligiel SEG, Voorhees JJ: All-*trans* retinoic acid and extracellular  $\text{Ca}^{2+}$  differentially influence extracellular matrix production by human skin in organ culture. *Am J Pathol* 1993, 142:1813-1822
- Angeles AM, Kahari V-M, Chen YQ, Ramirez F, Uitto J: Enhanced collagen gene expression in fibroblast cultures treated with all-*trans* retinoic acid: evidence for up-regulation of the  $\alpha 2$  (1) promoter activity. *J Invest Dermatol* 1990, 94:540a
- Schwartz E, Cruikshank FA, Mezick JA, Kligman H: Topical all-*trans* retinoic acid stimulates collagen synthesis *in vivo*. *J Invest Dermatol* 1991, 96:976-978
- Clark SD, Kobayashi DK, Welgus HG: Regulation of the expression of tissue inhibitor of metalloproteinases and collagenase by retinoids and glucocorticoids in human fibroblasts. *J Clin Invest* 1987, 80:1280-1288
- Griffiths CEM, Russman A, Majmudar G, Singr RS, Hamilton TA, Voorhees JJ: Restoration of collagen formation in photodamaged human skin by tretinoin (retinoic acid). *N Engl J Med* 1993, 329:530-535
- Kligman LH: Effects of all-*trans* retinoic acid on the dermis of hairless mice. *J Am Acad Dermatol* 1986, 15:779-785
- Kligman LH, Duou CH, Kligman AM: Topical retinoic acid enhances the repair of ultraviolet damaged dermal connective tissue. *Connect Tissue Res* 1984, 12: 139-150
- Chen S, Kiss I, Trampusch KM: The priming effect of ultraviolet B radiation on retinoic acid-stimulated collagen synthesis in the mouse photo-damage model. *Photoderm Photoimmunol Photomed* 1992, 9:104-108
- Praeger FC, Cristofalo VJ: Modulation of WI-38 cell proliferation by elevated levels of  $\text{Ca}^{2+}$ . *J Cell Physiol* 1986, 129:27-35
- Praeger FC, Gilchrist BA: Influence of increased extracellular calcium concentration and donor age on density-dependent growth inhibition of human fibroblasts. *Proc Soc Exp Biol Med* 1986, 182:315-321
- Praeger FC, Cristofalo VJ: Effects of elevated levels of extracellular  $\text{Ca}^{++}$  on young and old WI-38 cells. *In Vitro* 1980, 16:239-249
- Praeger FC, Cristofalo VJ: Age-related loss of response to elevated  $[\text{Ca}^{2+}]$  by WI-38 cells. *Gerontologist* 1984, 24:226-227
- Boynton AL, Whitfield JF, Isaacs RJ, Morton HJ: Control of 3T3 cell proliferation by calcium. *In Vitro* 1974, 10: 12-17
- Boynton AL, Whitfield JF, Isaacs RJ, Tremblay R: The control of human WI-38 proliferation by extracellular calcium and its elimination by SV-40 virus-induced proliferative transformation. *J Cell Physiol* 1977, 92:241-248
- Varani J, Shayevitz J, Perry D, Mitra RS, Nickoloff BJ, Voorhees JJ: Retinoic acid stimulation of human dermal fibroblast proliferation is dependent on suboptimal extracellular  $\text{Ca}^{2+}$  concentration. *Am J Pathol* 1990, 136:1275-1281
- Varani J, Gendimenico GJ, Shah B, Gibbs D, Capetola RJ, Mezick JA, Voorhees JJ: A direct comparison of the

- pharmacological effects of retinoids on skin cells *in vitro* and *in vivo*. *Skin Pharmacol* 1991, 4:254–261
23. Phan SH, Varani J, Smith D: Rat lung fibroblast collagen metabolism in bleomycin-induced pulmonary fibrosis. *J Clin Invest* 1985, 76:241–247
24. Liu SC, Karasek M: Isolation and growth of adult human epidermal keratinocytes in cell culture. *J Invest Dermatol* 1978, 71:157–162
25. Grynkiewicz G, Poenie M, Tsien TY: A new generation of  $\text{Ca}^{2+}$  indicators with greatly improved fluorescence properties. *J Biol Chem* 1985, 260:3440–3450
26. Ozaki Y, Yatomi Y, Kume S: Evaluation of platelet calcium ion mobilization by the use of various divalent ions. *Cell Calcium* 1992, 13:19–27
27. Kwan C-Y, Putney JW Jr: Uptake, intracellular sequestration of divalent cations in resting and methacholine-stimulated mouse lacrimal acinar cells. *J Biol Chem* 1990, 265:678–684
28. Hughes AD, Schachter M: Multiple pathways for entry of calcium and other divalent cations in a vascular smooth muscle cell line (A7r5). *Cell Calcium* 1994, 15:317–330
29. Betsholtz C, Westermarck B: Growth factor-induced proliferation of human fibroblasts in serum-free culture depends on cell density and extracellular calcium concentration. *J Cell Physiol* 1984, 118:203–210
30. Takemura H, Hughes AR, Thastrup O, Putney JW Jr: Activation of calcium entry by the tumor promoter thapsigargin in parotid acinar cells. *J Biol Chem* 1989, 264:12266–12271
31. Kass GEN, Llopis J, Chow SC, Duddy SK, Orrenius S: Receptor-operated calcium influx in rat hepatocytes. *J Biol Chem* 1990, 265:17486–17492
32. Jacob R: Agonist-stimulated divalent cation entry into single cultured human umbilical vein endothelial cells. *J Physiol* 1990, 421:55–77
33. Murray RK, Kotlikoff MI: Receptor-activated calcium influx in human airway smooth muscle cells. *J Physiol* 1991, 435:123–144
34. Berridge MJ: The biology, and medicine of calcium signalling. *Mol Cell Endocrinol* 1994, 98:119–124
35. Benham CD, Tsien RW: Calcium-permeable channels in vascular smooth muscle: voltage-activated, receptor-operated, and leak channels. *Soc Gen Physiol Ser* 1987, 42:45–64
36. Soltoff SP, Cantley LC: Mitogens and ion fluxes. *Annu Rev Physiol* 1988, 50:207–223
37. Estacion M: Characterization of ion channels seen in subconfluent human dermal fibroblasts. *J Physiol* 1991, 436:579–601
38. Chytil F, Sherman DR: How do retinoids work? *Dermatologica* 1987, 179:8–12
39. Darmon M, Rocher M, Cavey M-T, Martin B, Rabilloud T, Delescluse C, Shroot B: Biological activity of retinoids correlates with affinity for nuclear receptors but not for cytosolic binding protein. *Skin Pharmacol* 1988, 1:161–175
40. Petkovich M, Brand NJ, Krust A, Kastner P, Chambon P: A human retinoic acid receptor which belongs to the family of nuclear receptors. *Nature* 1987, 330:444–450
41. Meeks R, Zaharevitz GD, Chen RF: Membrane effects of retinoids: possible correlation with toxicity. *Arch Biochem Biophys* 1981, 207:141–147
42. Varani J, Inman DR, Perone P, Fligiel SEG, Voorhees JJ: Retinoid toxicity for fibroblasts and epithelial cells is separable from growth promoting activity. *J Invest Dermatol* 1993, 101:839–842
43. Astrom A, Petterson U, Krust A, Chambon P, Voorhees JJ: Retinoic acid and synthetic analogs differentially activate retinoic acid receptor-dependent transcription. *Biochem Biophys Res Commun* 1990, 173:339–345
44. Duell EA, Astrom A, Griffiths CEM, Chambon P, Voorhees JJ: Human skin levels of retinoic acid and cytochrome P-450-derived 4-hydroxyretinoic acid after topical application of retinoic acid *in vivo* compared to concentrations required to stimulate retinoic acid receptor-mediated transcription *in vitro*. *J Clin Invest* 1992, 90:1269–1274